

Natural scientific methods, a potential, or more needed in future archaeology?

DAGFINN MOE

From the earliest days of archaeology there has been continuous methodological development. Archaeological fieldwork today requires, as a minimum standard, stratigraphic and artefact description as well as collection of samples for independent dating, primarily ¹⁴C dating. Recent developments in fields dealing with the natural environment include many new methods applicable to palaeoenvironmental work that may also be of potential interest to those who deal with history and prehistory.

For some archaeologists it is an open question as to whether or not there is any need for cooperation with palaeoecologists. It is my opinion that such cooperation is not only important but very often necessary.

This paper deals firstly with different forms such cooperation may take, and secondly presents several recent palaeoenvironmental projects that have archaeological aspects, which at least for some researchers and for the public have provided new information about historic and prehistoric life ways.

Politicians as well as the general public today talk and ask about the environment in a broad sense, while most of the researchers today work within very narrow sectors or fields. For example, neither a botanist nor a geologist can manage to cover more than a small part within their field and be updated in the all the new techniques available.

To be able to communicate with the public, we specialists need firstly to communicate much more with each other, and secondly to learn how to answer questions from the public in general, including those from the media. Of special importance certainly is to answer, in a proper way, questions from the official institutions who normally support us financially.

Questions concerning prehistory from non-specialists most often have broad scope. The answers need to include information as to, when did it happen, what kind of climate existed, what kind of food-supply did the people have, what kind of small-game, big-game, fishes etc. existed at that time, what hunting strategies were used, and so on. During the last decades we have seen an increasing number of natural scientific methods in use in archaeological, – methods such as computers and laser equipment, different dating techniques, stratigraphic descriptions of sediment, chemical studies, as well as methods dealing with zoological and botanical fossils. Mineralogenic analysis might be of a special

interest in relation to studies of raw materials and ceramics to answer questions regarding artefact composition, origin, and tracing possible trade routes.

For most of us, the smaller archaeo-palaeoenvironmental projects, or the normal rescue excavation, are the most frequent ones, and a large number of them have been undertaken in Scandinavia (e.g. Hjelle et al. 1992; Olsen 1992). Some have been interdisciplinary projects, others multidisciplinary. There is a difference between these. The former normally involves cooperation between the participants from the beginning, continuing through fieldwork, and ends with a final discussion/presentation in a joint paper (Moe 1972; Moe et al. 1978). It is also within this joint team framework that unexpected questions or comments may stimulate unexpected new scientific ideas. Joint work, however, always need more time, and depends on mutual respect between the participants.

In multidisciplinary projects, the participants work separately (Moe 1972), and leave the project after having finished their paper or report. Most palaeoenvironmental projects are of this kind. They are easier to run, and the participants do not necessary need to agree on goals, or be involved in joint discussion. Misunderstandings are not always easy to avoid with regard to using each others terminology, suggestions or conclusions etc., and this can result in the publication of scientific errors.

A mixture of separate and joint papers is perhaps the best. At the same time each participant may present some work within their own field for their own scientific credit, parallel with new interesting joint chapters. This is important to remember and might be a problem in some cases.

Since 1992 many papers have been published on the findings of the Neolithic Ice Man in the Alps (Höphel et al. 1992; Spindler et al. 1995). A wide variety of techniques have been used, and the data available, at the moment is great (never has a body been studied better, and we know that still more material will be studied). Some of the technical methods used are well established and well known, but rarely used on prehistoric material. Others needed to be developed or adapted to the special problem or questions concern. Nearly everything regarding medical conditions, food, textiles, tools etc. will be known. As a botanist, it has certainly been of special interest to learn what plant, and tree species were used for the different tools that were found, which parts of the tree/shrub were used (Oeggli & Schoch 1995), as well as where these different species were collected.

Natural scientific measurements will always give some results. Depending on stratigraphy, sampling techniques, identification, contamination etc. the result(s) will fit into a model, or not. Some will sometimes try to hide/forget "wrong" data, while in many cases such data ought actually to receive extra attention - perhaps to the end of improving the technique used, and/or to modify a model. The "wrong" results therefore might be among the most stimulating ones scientifically.

It is not easy to extract a limited number of examples from the field of palaeoecological to demonstrate the effect of cooperation and joint work. References to some have already been made, and short comments on others follows.

An archaeological excavation at an iron age site above the tree line in South Norway, yielded a ¹⁴C date on burnt wood of around 8000 years BP. The archaeologist first became rather frustrated, however, the botanist provided a key to understanding the mystery after having identified some pieces of the charcoal left behind to pine (*Pinus sylvestris*). Fossil roots and stems from periods

when the tree line was higher are known from bogs at this altitude, and the wood dated at the iron age site obviously belonged to a former pine forest - from 8000 years ago (Moe & Odland 1992). Due to a lack of wood at the site in the Iron Age, the Iron Age man gathered an ancient pine stump from the local mire, dried it and used it for burning.

From the same area, fish bones of salt-water fish were found in an archaeological context in a high arctic-alpine region, that is, above the present tree line which is more than 1100 m.a.s.l. in South Norway (Indrelid 1994). Material of more or less the same age from the Medieval excavations at the old harbour in Bergen (Bryggen) included bones and antlers of reindeer, a species native to the high arctic-alpine area.

Behre and Oeggli's (1996) recent summary of archaeo-botanical results on early farming in Europe and adjacent areas is an important contribution to this kind of palaeo-environmental knowledge.

Some paths, trackways and trade routes are known from historic as well as prehistoric times (Moe & van der Knaap 1990). Coins are perhaps the best artefact for identifying trade, while jewelry, some kinds of ceramics, flint etc. may be more difficult to trace. Barrels, 1000 years old made of fir (*Abies alba*) were found in Haiteby/Hedeby in north Germany, close to the Danish border (Behre 1983). These demonstrate a relatively long distance transport of the barrels from south Germany, where fir grew, to the Danish border, where the content of the barrels, perhaps wine, was used.

A very detailed and interesting recent analysis of sheep/goat coprolites (faeces) in an archaeological context from Switzerland demonstrate use of twigs and branches of hazel (*Corylus avellana*), birch (*Betula* sp.) and alder (*Alnus incana*) as fodder. In addition the results show that this Neolithic settlement was occupied during winter and spring. (Rasmussen, 1991, 1993: Haas & Rasmussen 1992).

A pilot study of a human coprolite from Birka, central Sweden, shows the use of lime-honey (*Tilia*) as a sweetener in food (Hjelmtveit & Moe in prep). Perhaps the honey was used in the local mead? The origin of the lime-honey is still an open question, it may have been imported from South Sweden, or more likely from Poland or the Baltic republics.

In Norway both sea level and tree line changes are and have been very important for human occupation and general adaptation (Hafsten 1979; Moe et al. 1978). The sea level changes vary a great deal geographically, from a total uplift only of a few meters in the southwestern most part of the country, to more than 220 m around Oslo in East Norway. The techniques used to determine sea level changes are more or less standardised and normally involve stratigraphic studies and 14C dating together with pollen, diatom, macrofossil, microfossil and phytoplankton analyses. The changes might be of great interest to the understanding of topographic land mass changes, localisation of sounds with strong tidal currents, and for the understanding of changes in flora and fauna composition, perhaps sometimes caused by changing climate.

The first changes in tree line and forest limits took place in Late-Glacial/ Early Holocene time, during the main immigration (Berglund et al. 1995). After the climatic optimum in the Boreal period (Moe 1995) evidence for a lowering of the tree line is found in alpine and arctic-alpine areas in Europe. Major changes are found to have effected the big game hunting strategy during Mesolithic in

both the southern Norwegian arctic-alpine region (Moe et al. 1978) and most likely also in the Alps. (e.g. Broglio 1992; Fedele & Buzzetti 1993). Minor tree line changes are more difficult to trace but must have taken place everywhere and disturbed game-animal movement.

Natural changes in forest composition, for instance after the immigration of for example the shade trees as beech (*Fagus sylvatica*) and/or spruce (*Picea abies*), may easily have changed the natural conditions required by certain birds and mammals, and therefore also affected humans (Moe & Hjelle in prep.).

It is obvious that natural changes have occurred, and it is obvious that prehistoric man had to adapt the changes both at the coast, along the rivers and in the more inland areas to be able to survive.

Most of the various natural scientific methods used by archaeologist today are used and are being improved by separate and independent laboratories at colleges or universities. Most of them are in daily use as parts of accepted, specialised fields of zoology, botany or geology, and most educational institutions take care of their own training and development. Also international specialist journals such as "Vegetation History and Archaeobotany" and "Holocene" have been started during the last 5-6 years and address these fields. We do not need to be prophets to predict that within few decades, "environmentologists", as most of us are, will need to fight against a potential danger of isolation.

I challenge the archaeologist and the palaeoecologist to establish and maintain contact with each other. This will prevent any second class environmental studies among the archaeologists from resulting in a possible breakdown between themselves and the independant, ecological and palaeo-ecological scientific fields which are well established in many countries today. Further, the palaeoecologist needs knowledge and understanding of the presence of man, and of environmental use and disturbances, which in periods are stronger influences than natural changes. A natural interaction will take place between fields on a reciprocal basis in joint projects, supporting the establishment of horizontal, scientific communication. It is, therefore, a great challenge for both archaeologists and archaeo-natural scientist to know each other better in the future, if we are interested in developing in our separate fields.

References

- Behre, K.E. 1983. *Ernährung und Umwelt der wikingerzeitlichen Siedlung Haithabu. Die Ergebnisse der Untersuchungen der Pflanzenresten*. Wachholtz, Neumünster.
- Behre, K.-E. & K. Oeggl 1966. *Early farming in the old world. Recent advances in archaeobotanical research*. Springer Verlag, Berlin.
- Berglund, B.E., H.J.B. Birks, M. Ralska-Jasiewiczowa & H.E. Wright (eds.) 1995. *Palaeoecological events during the last 15000 years: Regional syntheses of palaeoecological studies of lakes and mires*. John Wiley & Sons, Chichester.
- Broglio, A. 1992. Mountain sites in the context of the North-East Italian Upper Palaeolithic and Mesolithic. *Preistoria Alpina*, Museo Tridentino di Scienze Naturali (Trento), Vol. 28, No. 1, 293-310.
- Fedele, F. & M. Buzzetti 1993. Pian dei Cavalli: Sui passi dei primi uomini nelle Alpi. *Museo della Valchiavenna*, (Valchiavenna) Vol. 2.
- Hafsten, U. 1979. Late and post-Weichselian shore level changes in South Norway. In *The Quaternary history of the North Sea* (Oele, E., R.T.E. Schuttenhelm & A.J. Wiggers

- (eds.) p. 45-59. *Acta Univ. Ups. Symp. Univ. Ups. Annum Quingentesimum Celebrantis*, Vol. 2, Uppsala.
- Hjelle, K.L., A.K. Hufthammer, P.E. Kaland, A.B. Olsen & E.C. Soltvedt 1992. *Kotedalen - en boplass gjennom 5000 år. Vol. 2, Naturvitenskapelige undersøkelser*. 150 p. University of Bergen, Historisk museum.
- Hjelmtveit, S.M. & D. Moe (in prep.). A preliminary study into a coprolite deposit at Birka, Central Sweden.
- Höpfel, F., W. Platzer & K. Spindler (eds.) 1992. *Der Mann im Eis*. Vol. I. Universität Innsbruck, No. 187, Innsbruck.
- Indrelid, S. 1994. Fangstfolk og bønder i fjellet. Bidrag til Hardangerviddas førhistorie 8500-2500 før nåtid. 344 p. *Universitetets Oldsaksamling Skr.* 17.
- Moe, D. 1972. Samarbeid eller tverrfaglig forskning. *Forskningsnytt* (Norwegian Research Council, Oslo) Vol. 3, 54-55.
- Moe, D. 1995. Climatic variations in western Norway during the last 13,000 years. A review. *Geologija*, (Lituania) (1994) Vol. 17, 15.
- Moe, D. & K.L. Hjelle (in prep.). Mesolithic man adaptation to changing alpine/subalpine vegetation in the Central and Eastern Alps. (Based on a vegetational historical study from Val Febbraro, Valle Spluga, Italy).
- Moe D. & A. Odland 1992. The influence of the temperature climate on the vertical distribution of *Alnus incana* (Betulaceae) through the Holocene in Norway. *Acta Botanica Fennica*, Vol. 144, 35-49.
- Moe, D. & W.O. van der Knaap 1990. Transhumance in mountain areas: Additional interpretation of three pollen diagrams from Norway, Portugal and Switzerland. *Pact journal*, Vol. 31, 91-104.
- Moe, D., S. Indrelid & O. Kjos-Hanssen 1978. A study of environment and early man in the southern Norwegian highlands. *Norwegian Archaeological Review*, Vol. 11, No. 2, 73-83.
- Oeggli, K. & W. Schoch 1995. Neolithic plant remains discovered together with a mummified corpse ("Homo tyrolensis") in the Tyrolean Alps. *Res archaeobotanicae* (Kroll, H. & Pasternak, R. (eds.)), 229-238. International workgroup for palaeoethnobotany, Proceedings 9th Symposium IWGP 1992. Oetker-Voges-Verlag, Kiel.
- Olsen, A.B. 1992. *Kotedalen - en boplass gjennom 5000 år. Vol. 1. Fangstbosetning og tidlig jordbruk i vestnorsk steinalder: nye funn og nye perspektiver*. 271 p. University of Bergen, Historisk museum.
- Rasmussen, P. 1991. Leaf foddering of livestock in the neolithic: Archaeobotanical evidence from Weier, Switzerland. *Journal of Danish Archaeology*, 1989, Vol. 8, 51-71.
- Rasmussen, P. 1993. Analysis of goat/sheep faeces from Egolzwil 3, Switzerland: evidence for branch and twig foddering of livestock in the neolithic. *Journal of Archaeological science*, Vol. 20, 479-502.
- Spindler, K., E. Rastbichler-Zissering, H. Wilfing, D. zur Nedden & H. Northdurfter (eds.) 1995. *Der Mann im Eis*. Vol. II, 320 p. Springer-Verlag, Wien.

Gamtos mokslų metodai: ar tai tik galimybė, ar ateities archeologijos būtinybė?

DAGFINN MOE

Santrauka

Gamtos mokslų metodai yra taikomi archeologiniame darbe, tiriant aplinkos modelius, ir ateityje jie, matyt, įgis dar didesnę svarbą. Šiame darbe kalbama apie du bendradarbiavimo būdus: daugiadisciplininį ir tarpdisciplininį. Taip pat pateikiami kai kurie tokių bendrų projektų rezultatai, paminint ir šalutinį jų poveikį.

*Dagfinn Moe
Department of Botany, University of Bergen
Alleg. 41, 5007 Bergen, Norway*